


High-protein breakfast prevents body fat gain in obese teens



This article originally appeared on Dr. Galland's website, PillAdvised.

Go for high-protein breakfast!

Some 60 percent of young people habitually skip breakfast up to four times a week.

Although health experts recommend breakfast as a strategy to reduce the chance of obesity, little research has examined if the actual type of breakfast consumed plays a significant role in one's health and weight management.



University of Missouri researchers compared the benefits of consuming a normal-protein breakfast to a high-protein breakfast and found the high-protein breakfast – which contained 35 grams of protein – prevented gains of body fat, reduced daily food intake and feelings of hunger, and stabilized glucose levels among overweight teens who would

normally skip breakfast.

Heather Leidy, Ph.D., an assistant professor in the Department of Nutrition and Exercise Physiology at the MU School of Medicine and lead author of the study, says the key to eating 35 grams of protein is to consume a combination of high-quality proteins including milk, eggs, lean meats and Greek yogurt.

“This study examined if the type of breakfast consumed can improve weight management in young people who habitually skip breakfast,” said Leidy. “Generally, people establish eating behaviors during their teen years. If teens are able to develop good eating habits now, such as eating breakfast, it’s likely to continue the rest of their lives.”

Leidy and her colleagues fed two groups of overweight teens who reported skipping breakfast between five and seven times a week either normal-protein breakfast meals or high-protein breakfast meals. A third group of teens continued to skip breakfast for 12 weeks.

“The group of teens who ate high-protein breakfasts reduced their daily food intake by 400 calories and lost body fat mass, while the groups who ate normal-protein breakfast or continued to skip breakfast gained additional body fat,” Leidy said. “These results show that when individuals eat a high-protein breakfast, they voluntarily consume less food the rest of the day. In addition, teens who ate high-protein breakfast had more stable glucose levels than the other groups.”

Leidy says large fluctuations in glucose levels are associated with an increased risk of Type 2 diabetes among young people, which can make health complications associated with weight gain more intense.

The normal-protein breakfast meal was milk and cereal and contained 13 grams of protein. The high-protein breakfast meals included eggs, dairy and lean pork that contained 35

grams of protein. Participants in the groups were instructed to report feelings of hunger and their daily intakes of food and beverages. Their body weight and body composition were measured at the beginning and end of the 12-week period. In addition, the participants wore a device that assessed minute-to-minute glucose levels throughout the day.

Reference:

“A pilot study examining the effects of consuming a high-protein vs normal-protein breakfast on free-living glycemic control in overweight/obese ‘breakfast skipping’ adolescents,” L B Bauer, L J Reynolds, S M Douglas, M L Kearney, H A Hoertel, R S Shafer, J P Thyfault and H J Leidy, *International Journal of Obesity* (1 June 2015) | doi:10.1038/ijo.2015.101