

# Ask Leyla: Kidney Stones and the microbiome



**Q:** I had a kidney stone for the first time at 52 years of age, and I was researching why this happened at this stage of my life. I read how a gut bacteria called *Oxalobacter formigenes* metabolizes oxalates in the gut, reducing the incidence of kidney stones.

How can I colonize my gut with this helpful bacteria?

**A:** Approximately 75 percent of all kidney stones are calcium

oxalate stones. Oxalate is a common component of many nutritious plant foods such as spinach, nuts, berries, oats, beans, chard, watercress, black tea and cocoa, to name a few. However, the propensity of these foods to raise urinary oxalate is dependent not just on oxalate content, but how efficiently it's absorbed.

It's true when they say good health begins in the gut because oxalate degradation requires an adequate amount of *Oxalobacter formigenes*. This beneficial bacteria helps reduce the amount of oxalate absorbed decreasing the risk of kidney stone formation. While colonization of *O. formigenes* begins in infancy, only 60 to 80 percent of adults continue to carry the bacteria. According to researchers, this may be due to the use—and overuse—of antibiotics.

To date, studies reveal that *O. formigenes* as well as other probiotic bacteria could be protective in those predisposed to kidney stones. One study reported an average decrease of 40 percent in urinary oxalate excretion in six subjects with kidney stones after four weeks of taking probiotics, while a second study of eleven healthy individuals also following a low-oxalate diet showed a decrease of 33 percent in urinary oxalate. The probiotic formulations in these studies included the following species:

*Lactobacillus acidophilus*

*Lactobacillus plantarum*

*Lactobacillus paracasei*

*Lactobacillus bulgaricus*

*Streptococcus thermophilus*

*Bifidobacterium infantis*

*Bifidobacterium longum*

Notice these probiotics are commonly found in yogurt and kefir as well as probiotic formulations, so it's fairly easy to reinoculate your gut to support *O. formigenes* growth.

RELATED: Ask Leyla: How much water should I be drinking every day?

There are other risk factors for kidney stone formation. Not drinking enough water is one. Generally, we need eight, eight-ounce cups a day. In fact, the highest incidence of kidney stones occurs in summer, and it's usually secondary to dehydration. Taking calcium supplements (especially without magnesium) or a high calcium diet as well as taking too much vitamin C will also contribute to stone formation.

If you're predisposed to kidney stones, taking a daily magnesium citrate supplement is recommended to help keep calcium from crystallizing and forming stones. Drinking water with lemon also introduces citrate in the form of citric acid, aiding in solubilizing stones.

To your health!

Liebman M, Al-Wahsh I. Probiotics and other key determinants of dietary oxalate absorption. *Amer Society for Nutr.* 2011; 2: 254-260.

Leyla Muedin, MS, RD, CDN

*Email your questions to [RadioProgram@aol.com](mailto:RadioProgram@aol.com).*